

Journal Pre-proof

Antidepressants' effects on testosterone and estrogens: what do we know?

Pavlina Pavlidi, Nikolaos Kokras, Christina Dalla



PII: S0014-2999(21)00151-5

DOI: <https://doi.org/10.1016/j.ejphar.2021.173998>

Reference: EJP 173998

To appear in: *European Journal of Pharmacology*

Received Date: 1 November 2020

Revised Date: 22 February 2021

Accepted Date: 1 March 2021

Please cite this article as: Pavlidi, P., Kokras, N., Dalla, C., Antidepressants' effects on testosterone and estrogens: what do we know?, *European Journal of Pharmacology*, <https://doi.org/10.1016/j.ejphar.2021.173998>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

Credit Author Statement

Pavlina Pavlidi: Writing - Original draft preparation

Christina Dalla and Nikolaos Kokras: Conceptualization, Reviewing and Editing

Antidepressants' effects on testosterone and estrogens: what do we know?

Pavlina Pavlidi¹, Nikolaos Kokras^{1,2*}, Christina Dalla^{1*}

Affiliations

1. Department of Pharmacology, Medical School, National and Kapodistrian University of Athens, Athens, Greece.
2. First Department of Psychiatry, Eginition Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece.

* These two authors contributed equally to this work

Corresponding Author:

Christina Dalla, Department of Pharmacology, Medical School, National and Kapodistrian University of Athens, Mikras Asias 75, 11527, Goudi, Athens, Greece, cdalla@med.uoa.gr

Abstract

Various antidepressants are commonly used to treat depression and anxiety disorders, and sex differences have been identified in their efficacy and side effects. Steroids, such as estrogens and testosterone, both in the periphery and locally in the brain, are regarded as important modulators of these sex differences. This review presents published data from preclinical and clinical studies that measure testosterone and estrogen level changes during and/or after acute or chronic administration of different antidepressants. The majority of studies show an interaction between sex hormones and antidepressants on sexual function and behavior, or in depressive symptom alleviation. However, most of the studies omit to investigate antidepressants' effects on circulating levels of gonadal hormones. From data reviewed herein, it is evident that most antidepressants can influence testosterone and estrogen levels. Still, the evidence is conflicting with some studies showing an increase, others decrease or no effect. Most studies are conducted in male animals or humans, underscoring the importance of considering sex as an important variable in such investigations, especially as depression and anxiety disorders are more common in women than men. Therefore, research is needed to elucidate the extent to which antidepressants can influence both peripheral and brain levels of testosterone and estrogens, in males and females, and whether this impacts the effectiveness or side effects of antidepressants.

Keywords

antidepressant; estrogens; testosterone; adverse effects; sexual dysfunction; hormones

1. Introduction

The pharmacotherapy of mood and anxiety disorders relies heavily on drugs that modulate the monoaminergic neurotransmission (Yohn et al., 2017). Currently, selective serotonin reuptake inhibitors (SSRIs), comprising of citalopram/escitalopram, sertraline, fluoxetine, paroxetine, and fluvoxamine, is the most widely used class of such medications (Pirraglia et al., 2003). Serotonin and Norepinephrine Reuptake Inhibitors (SNRIs), namely venlafaxine and duloxetine, is another widely used category of antidepressants. Their common primary mechanism of action is the inhibition of the presynaptic reuptake of serotonin and/or norepinephrine neurotransmitters (Artigas et al., 2002). The older generation of tricyclic antidepressants (TCAs), i.e., amitriptyline, imipramine, desipramine, nortriptyline, is less frequently used nowadays, as their non-selective blockade was associated with more side-effects (Pacher et al., 2001). Several other compounds, such as agomelatine, bupropion, mirtazapine, and trazodone are considered atypical antidepressants. Atypical antidepressants' mechanism involves modulation of dopamine neurotransmission, modulation of serotonin and melatonin receptors. Furthermore, recently esketamine was the first licensed antidepressant that acts on the NMDA receptors (Frazer, 1997; Garay et al., 2017).

Strong evidence supports the efficacy and safety of those medications in treating mainly depression and secondarily anxiety disorders. However, in neuroscience and neuropsychopharmacology, there is a gradually increasing research interest regarding sex differences in the symptomatology, prevalence, etiology, and treatment of psychiatric disorders. Indeed, sex differences have been described in mood and anxiety disorders, in cognitive processes as well as in physiological and especially neuroendocrine mechanisms that present considerable differences among men and women (Dalla et al., 2010; Kokras and Dalla, 2014, 2017). Nonetheless, the study of both sexes in trials investigating psychiatric

drugs is still problematic, potentially leading to inadequate evaluation of sex differences ([Kokras et al., 2019](#)). For instance, several studies have described gender differences in men and women's pharmacokinetic profiles for several antidepressants. In particular, body weight, volume of blood, enzyme activity, and clearance rates have been identified as potential sources of these differences ([Kokras et al., 2011](#); [Sramek et al., 2016](#)).

Moreover, many studies suggest a significant interaction between sex hormones and antidepressants ([Damoiseaux et al., 2014](#); [Pawluski et al., 2020](#); [Sramek and Cutler, 2011](#); [Westlund Tam and Parry, 2003](#); [Williams and Trainor, 2018](#)), while focusing only on the aftereffect of antidepressants' influences on sexual function and behavior. Unfortunately, the exploration of possible antidepressant-related changes in hormonal levels still remains significantly understudied. In particular, most studies focus on the possible synergistic effect of hormones and antidepressants, rather than investigating antidepressants' effects on circulating levels of male and female hormones, such as testosterone and estrogens.

Therefore, the purpose of this review, following an exhaustive literature search, is to present preclinical and clinical studies that evaluate changes in testosterone and estrogen levels during and/or after acute or chronic administration of different antidepressants. Furthermore, we aim to elucidate the extent to which antidepressants can influence both serum and brain levels of testosterone and estrogens, in both males and females. Prompted by the extended relations between depression and alterations in the hormonal milieu, it is of great importance to determine the effects of antidepressants on testosterone and estrogen levels. Therefore, the aim of this review is to explore the drugs' mechanisms of action and possibly further elucidate the pathophysiology of depression.

2. Testosterone

Testosterone is one of the major androgens biosynthesized during steroidogenesis from cholesterol (Figure 1) ([Hu et al., 2010](#)). Interestingly, some preclinical and clinical studies suggest that antidepressants can affect the hormonal milieu in both sexes, and data are presented herein.

2.1. Citalopram and escitalopram

A study conducted by [Ilgın et al.](#) in male rats investigated the chronic effects of citalopram administration in sexual function. The authors reported elevated serum testosterone levels in citalopram-treated animals, regardless of the administered dose ([Ilgın et al., 2017](#)). In accordance, [Przegaliński et al.](#) reported enhanced testosterone concentration in two brain areas (cerebral cortex, and hypothalamus) following citalopram administration ([Przegaliński et al., 1987](#)). On the contrary, adult male rats treated chronically with escitalopram exhibited reduced serum testosterone levels compared to controls ([Erdemir et al., 2014](#)). To date, no human clinical studies have explored the effects of citalopram and escitalopram on testosterone levels.

2.2. Sertraline

A recent study investigated the possible endocrine disruptions of sub-chronic sertraline administration on sex steroid production. Male rats were treated with 3 different doses of sertraline for 15 days and all doses caused an extensive endocrine disruption, both directly on steroid production in major endocrine tissues, and indirectly by affecting gene expression. In particular, it was found that sertraline administration caused a reduction in brain and testis levels of testosterone, an augmentation in adrenal levels, and had no effect in plasma testosterone levels ([Munkboel et al., 2018](#)). Similar results were reported in adult male rats, which received long-term treatment with sertraline and exhibited decreased

testosterone serum levels ([Erdemir et al., 2014](#)) ([Bezerra et al., 2019](#)). A recent study evaluated the reproductive aspects of sertraline on pubertal male rats, and testosterone levels were significantly reduced ([ElMazoudy et al., 2020](#)). Moreover, a study from our laboratory investigated the effects of sertraline on testosterone levels in adult male and female rats. Interestingly, subacute sertraline treatment decreased remarkably testosterone levels only in males ([Kokras et al., 2015](#)) (Figure 2). On the contrary, two studies evaluated the possible mechanisms that underlie chronic sertraline-induced reproductive toxicity in male rats, and both studies reported an increase in serum testosterone levels following sertraline administration ([Atli et al., 2017](#); [Hamdi, 2019](#)). A possible explanation for the contradiction in findings described above may be the differences in the studies' experimental design. More specifically, Kokras et al. examined the sertraline effects in adult rats after intraperitoneal administration of the antidepressant ([Kokras et al., 2015](#)). In comparison, the other two studies investigated sertraline-related changes during adolescence, while administering the antidepressant via oral gavage ([Atli et al., 2017](#); [Hamdi, 2019](#)). No human studies have focused on testosterone modulation by sertraline.

2.3. Paroxetine

Like the antidepressants mentioned above, paroxetine is another SSRI that has been shown to reduce testosterone serum levels in male rats by two recent studies ([Allouh et al., 2014](#); [Erdemir et al., 2014](#)) (Table 1). However, sub-chronic evaluation of paroxetine effects on nocturnal endocrinological profiles in healthy men indicated no significant changes in the testosterone levels, and the nocturnal testosterone secretion ([Schlosser et al., 2000](#)).

2.4. Fluoxetine

Several preclinical studies have evaluated the chronic effects of fluoxetine administration and resulted in contradictory results. A study by Mennigen et al. aimed to determine the effects of waterborne fluoxetine, including environmental concentrations, on the reproductive axis of sexually mature male goldfish ([Mennigen et al., 2017](#)). It was reported that fluoxetine exposure inhibited both basal and pheromone-stimulated testosterone levels in male goldfish ([Mennigen et al., 2010](#)). Another study concluded that chronic fluoxetine administration in male rats decreased testosterone serum concentrations ([Ayala et al., 2018](#)). Similarly, long-term fluoxetine administration in adult and young male rats caused a considerable decrease in testosterone levels, irrespective of the administration route ([Bataineh and Daradka, 2007](#); [Bezerra et al., 2019](#); [Sakr et al., 2015](#)). On the contrary, evaluation of sexually naive adult male rats' reproductive systems under chronic fluoxetine treatment indicated that serum testosterone levels were indistinguishable from controls ([Erdemir et al., 2014](#); [Taylor et al., 2004](#)). Similar to paroxetine, fluoxetine was found not to exert changes in serum testosterone levels in male outpatients with major depressive disorder ([Papakostas et al., 2006](#)). On the contrary, both placebo and fluoxetine administration for 1 month in male patients with depression and healthy controls resulted in considerable differences -both increases and decreases- in testosterone levels, indicating possible correlations with depression severity, apathy and treatment response ([Bell et al., 2006](#)).

2.5. Venlafaxine

Two studies in adult male rats concluded that venlafaxine increases testosterone levels and testosterone's aromatization to estrogens ([de Santi et al., 2020](#); [Eid et al., 2019](#)). Two case reports associated low testosterone levels in two male patients with venlafaxine treatment ([Bell and Shipman, 2000](#); [Shebak and Varma, 2014](#)). Both reports concluded that the onset of low testosterone levels during venlafaxine treatment was associated with apathy,

fatigue and reduced libido. Following venlafaxine discontinuation, testosterone levels were restored, suggesting that venlafaxine may affect sexual functioning by affecting testosterone. It is worth mentioning that both studies underlined the shortage of literature associating venlafaxine administration and testosterone levels, suggesting that this shortage is due to the rarity of this side effect, or due to the lack of monitoring of testosterone levels in patients with such symptoms (Bell and Shipman, 2000; Shebak and Varma, 2014). On the contrary, short-term use of venlafaxine in 21 non-depressed male patients with premature ejaculation did not change the testosterone levels (Kilic et al., 2005).

2.6. Duloxetine

Duloxetine has been sparsely studied with regards to its effects on testosterone. An *in vitro* study reported that duloxetine can cause endocrine disruption, with CYP17 enzyme being the primary target. The cytochrome CYP17 (P450 17) is a key enzyme located upstream in the steroidogenic pathway, and most steroid synthesizing tissue is depended on CYP17 function (Islin et al., 2018). No *in vivo* preclinical studies and no clinical studies have been published regarding duloxetine's effects on testosterone.

2.7. Amitriptyline

Previous studies have reported conflicting findings regarding the effects of amitriptyline on testosterone levels. A study investigating the effects of sub-chronic administration of amitriptyline reported reduced testosterone levels in male rats' hypothalamus (Przegalinski et al., 1987). On the other hand, chronic administration of amitriptyline in virgin adult female rats did not cause any changes on testosterone levels (Li et al., 2019). Steiger et al. investigated testosterone's nocturnal secretion in male patients with major depression, before treatment and after recovery and drug cessation. It was shown that

an oral amitriptyline dose of 150 mg/day increased testosterone concentration after remission (Steiger et al., 1991).

2.8. Imipramine

A study investigated the effects of repeated antidepressant administration on testosterone serum and brain concentration in rats. An oral chronic treatment with imipramine reduced testosterone levels in serum and the cerebral cortex (Przegalinski et al., 1987). Twenty-eight days of imipramine treatment in male inpatients with depression caused a tendency to increase nocturnal plasma testosterone concentration (Sonntag et al., 1996).

2.9. Desipramine

N-Monoalkyl substituted tricyclic antidepressants like desipramine are known to undergo cytochrome (P450) -mediated biotransformation in the liver to produce inhibitory metabolite-intermediate complexes with the enzyme. An *in vitro* study indicated that inhibition of testosterone hydroxylation pathways was enhanced by prior incubation of desipramine with NADPH and rat liver microsomes (McNeil and Murray, 1996). There were no *in vivo* preclinical and clinical studies regarding desipramine's effects on testosterone.

2.10. Bupropion

A recent study investigated the effect of postnatal chronic exposure to bupropion on puberty onset, reproductive results, and testosterone levels in male rats, and reported a significant increase in testosterone levels, suggesting that this may affect puberty onset (Yardimci et al., 2019). Increased testosterone levels were also reported by another study after bupropion administration in male Wistar rats (Kankash et al., 2014). Studies suggest that the beneficiary effects of bupropion in sexual function and desire are not due to the

drug's influence on the hormonal milieu of men or women (Abdel-Hamid and Saleh el, 2011; Ishak et al., 2010; Miller et al., 2018; Patel et al., 2016). Despite that, a placebo-controlled trial of bupropion as a treatment for SSRI-induced sexual dysfunction, reported that desire/frequency showed significantly greater improvement in patients from both genders. Furthermore, the frequency was significantly correlated to total testosterone levels at baseline and week 4 of bupropion administration in women (Clayton et al., 2004).

2.11. Trazodone

Ilgin et al. identified elevated serum testosterone levels after trazodone administration in male rats, accompanied by oxidative stress and testicular damage (Ilgin et al., 2018). On the other hand, chronic trazodone administration in adult male rats reduced serum testosterone levels (Ashraf et al., 2014). Trazodone has been reported to cause priapism and enhanced libido while prolonging nocturnal erections (Meinhardt et al., 1999; Pyke, 2020; Vitezic and Pelcic, 2002). However, it is still unclear whether trazodone's effectiveness against erectile dysfunction is accomplished by changes in serum testosterone levels. It was shown that trazodone has a similar success rate as testosterone when used as a treatment towards erectile dysfunction. However, no serum testosterone levels were measured after treatment (Aydin et al., 1996). No clinical studies have examined the effects of trazodone on testosterone.

2.12. Esketamine

Ketamine is used as an anesthetic and acts as a noncompetitive antagonist of N-methyl-D-aspartate-type glutamate receptors. A nasal spray of low doses of esketamine was recently approved as a rapidly acting antidepressant for treatment-resistant depression (Bahr et al., 2019). A study utilizing wild-type (WT) zebrafish (*Danio rerio*) embryos demonstrated

that ketamine significantly elevated testosterone levels in comparison to controls (Trickler et al., 2014). Furthermore, Nazian et al. tested the acute effects of ketamine as an anesthetic agent in immature and young adult male rats that were either castrated or unoperated. Serum testosterone levels declined in intact immature male rats in response to ketamine, whereas castrated immature male rats responded to ketamine with significant serum testosterone increases (Nazian, 1988). Comparable results were also obtained by two studies, suggesting that chronic ketamine administration, regardless of dose, can lead to decreased serum concentrations of testosterone (El Shehaby et al., 2020; Qi et al., 2017). On the contrary, Gould investigated the effect of ketamine on plasma testosterone concentration in male sexually naïve rats and concluded that the combination of ketamine with xylazine followed by CO₂ euthanasia does not affect testosterone concentrations (Gould, 2008). Lastly, suppressed serum testosterone levels were also reported by a study utilizing adult female rats (Lee et al., 2000). Few studies have focused on investigating the ketamine effects on testosterone levels in nonhuman primates. For instance, a study performed in male nonhuman primates, compared metabolic clearance rates, production rates, and serum levels of testosterone in conscious and ketamine anesthetized adult male rhesus monkeys. A significant increase in metabolic clearance rate after ketamine anesthesia was observed. Serum testosterone levels and production rates were not significantly different between conscious and ketamine anesthetized animals (Zaidi et al., 1982). Similarly, single or multiple injections and chronic administration of ketamine hydrochloride in adult male rhesus monkeys and male cynomolgus monkeys caused no effects on serum testosterone levels (Malaivijitnond et al., 1998; Puri et al., 1981). Interestingly, only one study by Rodas-Martínez et al. reported that following capture, restraint, and anesthesia with ketamine (every 2 months for a year), adult male Spider monkeys exhibited a reduction in testosterone levels (Rodas-Martinez et al., 2013). It is worth mentioning that ketamine has been suggested to

decrease Luteinizing Hormone-Releasing Hormone (LHRH) level one hour after administration, and since LHRH and testosterone operate in a feedback mechanism, it is difficult to ascertain whether ketamine/xylazine exert an effect on LHRH, testosterone or both (Gould, 2008). No studies to date have reported measurements in testosterone levels following esketamine as an antidepressant treatment. However, a clinical study with twenty-seven male patients who received ketamine anesthesia (2 mg/kg, i.v.), reported decreased plasma concentration of testosterone during and after ketamine administration. The lowest concentration was detected on the first day after surgery (Oyama et al., 1977).

3. Estrogens

Estrogen is a major reproductive hormone, responsible mostly for the female reproductive system. Aromatase (CYP19A) is the key enzyme in the estrogens' biosynthesis, acting via androgens' aromatization. There are three major endogenous estrogens with estrogenic hormonal activity: estrone (E1), estradiol (E2), and estriol (E3) (Figure 1). Both men and women have lower levels of circulating estrogens compared to androgens, and although males have significantly lower levels of estrogens than females, they have important physiological roles in both sexes. Both preclinical and clinical studies have supported estrogens' involvement in the pathogenesis of depression and the augmentation of antidepressants' effectiveness (Berlanga and Flores-Ramos, 2006; Bryant et al., 2006; Sramek et al., 2016). For instance, regarding the pathogenesis of depression, most prior research has described that estradiol decline is associated with depression and anxiety in postmenopausal women (Bromberger and Epperson, 2018; Freeman et al., 2006; Ryan et al., 2009; Soares et al., 2001).

3.1. Citalopram and escitalopram

Preclinical studies have described that citalopram and escitalopram do not influence estradiol concentration in serum or the brain. Similarly, escitalopram does not share the estrogenicity seen in other SSRIs, i.e., fluoxetine (Montagnini et al., 2013; Przegalinski et al., 1987). Estrogenicity or estrogenic activity is defined as when compounds, such as antidepressants, mimic human estrogen by binding to estrogen receptors (ER), ER α and/or ER β . However, chronic citalopram administration (15 weeks) exerted hormonal effects in nonhuman primates, via elevating estrogens' secretion in stress-sensitive animals, but not in stress-resilient monkeys (Bethea et al., 2011; Weissheimer et al., 2010).

3.2. Sertraline

Uterus weight has been used as an indication of estrogens in female rats. Sertraline has been administered in both male and female and did not alter females' uterus weight nor affected estrogen levels in males and females (Montagnini et al., 2013). However, a tendency for a sertraline-induced dose-dependent reduction of estrogen levels was previously reported (Kokras et al., 2015). Interestingly, no human studies have focused on sertraline's effects on estrogens.

3.3. Fluoxetine

Lupu et al assessed whether the active metabolite of fluoxetine, norfluoxetine, interacts *in vitro* with the ER α and ER β . It was reported that norfluoxetine could induce estrogenic effects and potentiate estradiol's activity *in vitro* (Lupu et al., 2015). Similarly, it was shown that different doses of fluoxetine could increase estradiol levels *in vitro*, in a human adrenocortical carcinoma cell line (H295R), without affecting the aromatase mRNA expression (Lupu et al., 2017). Furthermore, another *in vitro* study showed that fluoxetine could downregulate ER α and ER β (Lupu et al., 2018). Fluoxetine exposure of male goldfish

led to the disruption of the reproductive endocrine axis, via increasing estrogen levels, the ER α , CYP19A1 enzyme, thus inducing estrogen-like effects in males ([Mennigen et al., 2017](#); [Silva de Assis et al., 2013](#)). On the contrary, studies investigating the effects of fluoxetine in female goldfish and sexually mature zebrafish showed a significant decrease of circulating E2, and conversely, a significant increase in circulating Luteinizing hormone (LH) and ovarian aromatase mRNA levels, suggesting disruption of E2-mediated feedback on LH release ([Lister et al., 2009](#); [Mennigen et al., 2008](#)). In the same teleost model, fluoxetine administration reduced ER β mRNA expression in both the hypothalamus and the telencephalon. However, fluoxetine exposure did not affect aromatase levels ([Mennigen et al., 2008](#)). On the contrary, short-term (96h) waterborne fluoxetine exposure of female Murray-Darling rainbowfish caused downregulation of aromatase mRNA and upregulation of ER α mRNA levels. However, concentrations of E2 in plasma were significantly lower than controls in response to fluoxetine, which could be influenced by decreased estrogen biosynthesis ([Bain et al., 2016](#)). Interestingly, a recent study reported that fluoxetine had no impact on E2 levels in female rainbow trout ([Monson et al., 2019](#)), whereas a study conducted in both sexes of zebra mussels showed a significant increase of endogenous levels of the esterified estradiol ([Lazzara et al., 2012](#)). Contrary to teleost studies, investigations of fluoxetine's influences on hormonal levels in rodents, and nonhuman primates have provided conflicting evidence. A study utilizing female adult rats reported that serum estradiol levels were not significantly different in rats injected with vehicle or fluoxetine ([Uphouse et al., 2006](#)). Similar results were reported by another study, which concluded that acute treatment with fluoxetine could inhibit lordosis behavior in female rats without affecting the gonadal hormones ([Miryala et al., 2013](#)). On the contrary, Taylor et al. reported that fluoxetine administration, regardless of dose, is sufficient to lower circulating estrogen titers ([Taylor et al., 2004](#)). A clinical study performed in normally cycling women suffering from major

depression and age-matched normally cycling non-depressed women evaluated the association of depression with changes in diurnal rhythms of free estradiol, in different phases of the menstrual cycle. Patients suffering from depression received fluoxetine treatment for 3 months and showed significantly higher peaks, lower troughs, and consequently, larger amplitudes of free estradiol than controls. However, these findings were not entirely attributed to fluoxetine treatment, but rather to the combination of depression, treatment, and menstrual phases. The authors concluded that further studies are required to differentiate the effects of fluoxetine on free estradiol (Bao et al., 2004).

3.4. Fluvoxamine

An *in vitro* study reported that human cytochrome P450 was the most active in catalyzing 2- and 4-hydroxylations of estradiol and estrone, and fluvoxamine exposure efficiently inhibited the estrogen hydroxylation in human liver samples (Yamazaki et al., 1998). No *in vivo* preclinical and clinical studies have focused on fluvoxamine's effects on estrogens.

3.5. Venlafaxine

A recent study evaluated the influence of chronic venlafaxine administration on testicular and sperm parameters, utilizing male Holtzman rats. It was shown that along with impaired sperm morphology venlafaxine decreased aromatase, and estrogen levels (de Santi et al., 2020). A preclinical study reported that venlafaxine administration could alleviate depressive symptoms of adult ovariectomized (OVX) female Wistar rats, and in parallel, increase serum estradiol levels (Saad et al., 2019). No clinical studies have focused on venlafaxine's effects on estrogens.

3.6. Amitriptyline

Regarding amitriptyline's effects on hormonal status, one study reported that repeated oral treatment with 10 mg/kg of amitriptyline did not influence estradiol concentration in serum or in the brain of male Wistar rats (Przegalinski et al., 1987). Furthermore, it was shown that amitriptyline could improve ovarian morphology and restore estradiol levels in female rats with induced Polycystic Ovary Syndrome (Li et al., 2019). No clinical studies were conducted investigating the effects of amitriptyline on estrogen levels.

3.7. Imipramine

Interestingly, an *in vitro* study utilizing CV-1 cells (monkey kidney) and C6-cells (rat glioma) concluded that imipramine directly activates the ER α (Hermann et al., 2001). However, no further investigation on estrogen levels was conducted. Results reported by preclinical studies suggested that chronic oral administration of imipramine was not sufficient to influence estradiol concentration in serum or in the brain of male Wistar rats (Przegalinski et al., 1987). No clinical studies explored imipramine's results in estrogen levels.

3.8. Desipramine

Similarly to imipramine, neither desipramine activated the ER α in an *in vitro* study utilizing CV-1 cells (monkey kidney) and C6-cells (rat glioma) (Hermann et al., 2001). Our literature research resulted in no *in vivo* preclinical and clinical studies investigating the effects of desipramine on hormonal levels.

3.9. Agomelatine

A recent study reported that sub-chronic agomelatine administration in young adult OVX female rats upregulated ER α and ER β mRNA expression in the hippocampus and

overall elevated serum estradiol levels (Okano et al., 1988). No clinical studies included the examination of agomelatine's effects on estrogen levels.

3.10. Bupropion

Clinical studies have associated low estrogen levels with decreased sexual desire in women. Reduced sexual drive is often treated with bupropion, but no studies have examined bupropion's possible influence on estrogen levels (Fooladi and Davis, 2012; Kingsberg and Woodard, 2015). No *in vivo* preclinical and clinical studies were found.

3.11. Esketamine

Ketamine significantly decreased E2 levels in zebrafish (*Danio rerio*) embryos (Trickler et al., 2014). Concordant data from mammalian studies suggest that ketamine attenuates ovary CYP19A1A aromatase expression but not the brain form of aromatase (CYP19A1B) in ketamine-treated embryos (Trickler et al., 2014). To that extend, a study by Lee et al. concluded that in female rats ketamine significantly decreased estradiol serum levels (Lee et al., 2000). On the contrary, the association between ketamine and decreased estradiol levels was not found in another study that measured rat uterine ER α and ER β after ketamine anesthesia (Zarembka et al., 1989). Ketamine effect on estrogen serum levels was also not significant in nonhuman primates (Bonney et al., 1979; Channing et al., 1977; Hutz et al., 1988). Only one clinical study measured estradiol levels after administering ketamine as an anesthetic in healthy women but found no effect (Sterzik et al., 1994).

4. Discussion

In this review, we presented published data on antidepressants' effects on testosterone and estrogen levels in males and females. In tables 1 to 4 we present all effects, i.e., the effect

of an antidepressant on testosterone (Table 1 and 2) or estrogens (Table 3 and 4) levels, from research papers and case reports, while including important information, such as sex, strain, age, and administered dose. Interestingly, many other reports mainly focus on the aftereffects of antidepressant's influences on sexual function and behavior, whereas they do not measure hormone levels. Nevertheless, the data presented herein show that antidepressants can influence the circulating levels of testosterone and estrogens. However, a precise conclusion regarding each antidepressant's effect, i.e., increase, decrease, and no effect, on hormonal levels cannot be drawn, as many studies report conflicting results.

This discrepancy in the findings may be due to strain, age, dose, route and duration of administration, and more importantly, sex differences. Differences may also arise from the evaluation of hormonal changes with various experimental techniques. Changes in the hormones' secretion's diurnal rhythms and the pulsative pattern of secretion may also explain why some studies fail to corroborate previous results when single time point measurements are performed. This concept and the arising methodological problems are well-identified for the Hypothalamic–pituitary–adrenal (HPA) axis and depression ([Kokras and Dalla, 2014](#)), but probably also applies to Hypothalamic–pituitary–gonadal (HPG) axis and its modulation in depression and antidepressant treatment ([Bao et al., 2004](#)). Studies in animals and primates suggest that HPA axis dysregulation and stress reactivity are present in depression, and antidepressants can facilitate normalization of the HPA axis function. Therefore, given that the HPA and HPG axis functions interact, there may be a common neuroendocrine response to antidepressant treatment ([Steiger et al., 1991](#)). However, the effect of antidepressants on the central regulation of the HPG axis still remains an unexplored issue. Although changes in peripheral levels of estrogens and testosterone have been reported, there is a considerable lack of quality evidence regarding their effects on LH and Follicle-Stimulating Hormone (FSH) levels. Interestingly, Kilic et al. and Oyama et al. reported that venlafaxine increased

FSH and ketamine increased LH, respectively. Such findings may reinforce significant sex differences, considering that the central HPG axis regulation is immensely different between sexes ([Kilic et al., 2005](#); [Oyama et al., 1977](#)).

Furthermore, only a few of the reported clinical studies performed pre-treatment measurements of hormonal levels. Patients with depression may show varied alterations of hormones, and depending on whether hormones are affected or not, the net effects of antidepressant treatment may be hard to estimate, suggesting that a controlled environment and the ability to measure pre- and post-treatment levels, similar to preclinical studies, are of significant importance to address this issue and explore antidepressant effects ([Bell and Shipman, 2000](#)). Of note, large and significant inter-individual differences were noted in clinical studies with patients presenting from a 100% increase to a 40% decrease in testosterone levels after treatment. These inter-individual differences may conceal antidepressants' effects and result in the absence of a total overall effect. However, at an individual level, patients' vulnerability to severe hormonal fluctuations, as a result of antidepressants' treatment, may be more frequently detectable. Such hormonal fluctuations may cause the appearance of somatic changes, i.e., sexual dysfunction, psychological side-effects, or testosterone-related apathy ([Bell et al., 2006](#)).

In preclinical research, several *in vivo* studies have underlined the importance of estrogens and testosterone by investigating the effects of gonadectomy in males and females or by temporally and locally increasing or decreasing the levels of these hormones. Gonadectomy in animals is a commonly used way to control endogenous hormonal circulation, starting with a profound reduction of testosterone and estrogen/progesterone levels, in males and females, respectively. Overall, in most cases, ovariectomy in female rats and mice has been shown to induce anxiety and depressive-like behaviors ([Puga-Olguin et al., 2019](#)) ([Kokras et al., 2018](#); [Kokras et al., 2017](#)), but it is important to remember that

ovariectomy in animals simulates surgical menopause in women and not physiological menopause per se ([Puga-Olguin et al., 2019](#)). However, notably, the peripheral circulation of gonadal steroids can pass the blood-brain barrier and influence neurosteroids' dynamics in the brain ([Brandt and Rune, 2020](#)).

Therefore, although a considerable amount of preclinical and clinical studies has proposed associations between changes in the hormonal milieu and behavior and even the endocrine system, only a few studies have investigated the effects of treatments on hormonal levels. Regarding preclinical studies, although a much greater proportion of publications investigate the bidirectional interaction between hormones and antidepressants, still the beneficiary and supporting properties of hormones to antidepressant treatment remain of paramount importance. Most of these studies describe the importance of local production of neurosteroids in the brain, in mood regulation ([Kokras et al., 2012](#); [Kokras et al., 2018](#)), learning and memory ([Taxier et al., 2020](#)), stress and anxiety ([Zheng et al., 2020](#)), and neuroplasticity ([Kumar and Foster, 2020](#); [Sellers et al., 2015](#)), without considering that future investigations should measure the levels of neurosteroids during neurological and neuropsychiatric conditions. Similarly, several clinical studies have expressed the importance of ample circulating hormonal levels in patients receiving certain antidepressant treatment, while omitting to measure hormonal fluctuations prior- and post-treatment or investigate whether hormonal changes are involved in the mechanism of action of the antidepressants.

Based on the presented findings, certain cautious comparisons between drug classes and differences between acute and chronic drug administration can be made. Such conclusions should be drawn while bearing in mind the significant methodological differences in published studies that were mentioned before, and the limited number of studies systematically investigating the antidepressants' effects on hormones.

Firstly, according to the reviewed data from preclinical studies, SSRIs affect to a greater extent, both testosterone and estrogen serum levels compared to the rest of drug classes (Table 1, 3). In particular, more than 50% of the reviewed publications report changes in testosterone and estrogen levels after SSRI administration. The same conclusion cannot be drawn from the comparatively fewer studies that investigated other classes of monoaminergic antidepressants. Secondly, data indicate differences between acute and sub-chronic or chronic drug administration on testosterone and estrogen levels. On the one hand, acute antidepressant treatment either decreases or does not affect testosterone and estrogen levels. On the other hand, data from sub-chronic and chronic antidepressant treatment are conflicted, probably due to variable treatment duration and differences in the time and method of sampling.

Furthermore, from our reviewed data it appears that testosterone levels are more frequently affected by antidepressants in comparison to estrogen. More specifically, the majority of studies found no changes in estrogen levels following drug administration, whereas the rest of the studies reported either increased or decreased levels of testosterone in both males and females (Table 1, 3). Unfortunately, inconsistencies in methods, i.e., inclusion of both sexes, doses, age, duration, and strain, as well as the technical difficulties in measuring low and variable estrogen levels account for the conflicting data and impede any firm conclusions.

Several *in vitro* studies investigate the relation between SSRIs and production and circulation of estrogen and androgen levels ([Hudon Thibeault et al., 2019](#); [Lupu et al., 2017](#)). Serotonin has been shown to regulate the enzyme aromatase (CYP19), which is the biosynthetic enzyme of estrogens ([Hudon Thibeault et al., 2019](#)). Furthermore, molecular simulations indicate that some SSRIs have a significant affinity for the CYP19. More specifically, *in vitro* data suggest that sertraline and paroxetine administration induce

aromatase activity, whereas fluoxetine exert the opposite effect on aromatase activity (Hudon Thibeault et al., 2019). Similar results were recently obtained in an *in vitro* study, where paroxetine administration in healthy mice caused CYP19 suppression, and eventually decreased circulating estradiol levels. (Zha et al., 2017) Therefore, a possible mechanism that SSRIs affect circulating estrogen and androgen levels might be via interfering with aromatase CYP19 (Hansen et al., 2017).

Another possible mechanism of how antidepressants modify hormone levels might involve estrogen receptors. Fluoxetine has been shown to induce estrogenic effects and potentiate estradiol's activity via its interactions and high affinity with ER α and ER β (Lupu et al., 2015). Similarly, a preclinical *in vivo* study suggested that agomelatine administration in rats upregulates ER α and ER β mRNA expression in the hippocampus and elevates serum estradiol levels (Okano et al., 1988). Oddly, other *in vitro* and *in vivo* investigations conclude in downregulation of ER α and ER β or reduction of ER β mRNA expression after fluoxetine administration (Lupu et al., 2018; Mennigen et al., 2008). Therefore, more research is warranted on the interaction of antidepressants with estrogen receptors and the resulting effect on hormone levels.

Notably, it is evident that most of the studies examining the antidepressants' effects on testosterone levels mainly choose male animals or patients, disregarding entirely female subjects. Similar sex bias is also present in studies measuring estrogens' fluctuation after antidepressant administration. Last but not least, only a few studies include both sexes in their experimental designs. Hence, there is a great need to consider sex and gender as an important biological variable in preclinical and clinical research, focusing on understanding depression and its treatments (Gururajan et al., 2019; Shansky and Woolley, 2016); especially since the prevalence and symptomatology of mood disorders are significantly influenced by sex and gender (Dalla et al., 2010; Kokras and Dalla, 2014, 2017).

Conflict of Interest

NK and CD have received honoraria and financial support from Janssen-Cilag, Elpen S.A. and Medochemie S.A. None of those is relevant to this study.

Acknowledgements

This work has been supported by Hellenic Foundation for Research and Innovation (HFRI).

Grant Number: HFRI-FM17-1676.

References

- Abdel-Hamid, I.A., Saleh el, S., 2011. Primary lifelong delayed ejaculation: characteristics and response to bupropion. *J Sex Med* 8, 1772-1779.
- Allouh, M.Z., Daradka, H.M., Al Barbarawi, M.M., Mustafa, A.G., 2014. Fresh onion juice enhanced copulatory behavior in male rats with and without paroxetine-induced sexual dysfunction. *Exp Biol Med (Maywood)* 239, 177-182.
- Artigas, F., Nutt, D.J., Shelton, R., 2002. Mechanism of action of antidepressants. *Psychopharmacology bulletin* 36, 123-132.
- Ashraf, S., Shariati, M., Zamanpor, M., 2014. Effect of trazodone HCL on levels of gonadotropin hormone, testosterone and histological changes of testis in adult rat. *Journal of Applied Environmental and Biological Sciences* 4, 80-84.
- Atli, O., Baysal, M., Aydogan-Kilic, G., Kilic, V., Ucarcan, S., Karaduman, B., Ilgin, S., 2017. Sertraline-induced reproductive toxicity in male rats: evaluation of possible underlying mechanisms. *Asian J Androl* 19, 672-679.
- Ayala, M.E., Gonzales, A., Olivarez, R.M., Aragon-Martinez, A., 2018. Fluoxetine treatment of prepubertal male rats uniformly diminishes sex hormone levels and, in a subpopulation of animals, negatively affects sperm quality. *Reprod Fertil Dev* 30, 1329-1341.
- Aydin, S., Odabas, O., Ercan, M., Kara, H., Agargun, M.Y., 1996. Efficacy of testosterone, trazodone and hypnotic suggestion in the treatment of non-organic male sexual dysfunction. *Br J Urol* 77, 256-260.
- Bahr, R., Lopez, A., Rey, J.A., 2019. Intranasal Esketamine (Spravato(TM)) for Use in Treatment-Resistant Depression In Conjunction With an Oral Antidepressant. *P T* 44, 340-375.
- Bain, P.A., Basheer, V.S., Gregg, A., Jena, J.K., Kumar, A., 2016. In vitro nuclear receptor activity and in vivo gene expression analysis in Murray-Darling rainbowfish (*Melanotaenia fluviatilis*) after short-term exposure to fluoxetine. *Comp Biochem Physiol C Toxicol Pharmacol* 188, 1-8.
- Bao, A.M., Ji, Y.F., Van Someren, E.J., Hofman, M.A., Liu, R.Y., Zhou, J.N., 2004. Diurnal rhythms of free estradiol and cortisol during the normal menstrual cycle in women with major depression. *Horm Behav* 45, 93-102.
- Bataineh, H.N., Daradka, T., 2007. Effects of long-term use of fluoxetine on fertility parameters in adult male rats. *Neuro Endocrinol Lett* 28, 321-325.

- Bell, S., Shipman, M., 2000. Reduced testosterone level in a venlafaxine treated patient. *Ann Clin Psychiatry* 12, 171-173.
- Bell, S., Shipman, M., Bystritsky, A., Haifley, T., 2006. Fluoxetine treatment and testosterone levels. *Ann Clin Psychiatry* 18, 19-22.
- Berlanga, C., Flores-Ramos, M., 2006. Different gender response to serotonergic and noradrenergic antidepressants. A comparative study of the efficacy of citalopram and reboxetine. *J Affect Disord* 95, 119-123.
- Bethea, C.L., Lima, F.B., Centeno, M.L., Weissheimer, K.V., Senashova, O., Reddy, A.P., Cameron, J.L., 2011. Effects of citalopram on serotonin and CRF systems in the midbrain of primates with differences in stress sensitivity. *J Chem Neuroanat* 41, 200-218.
- Bezerra, M.S., Martins, A.B.M., Trajano, F.M.G., Pontes, T.H.A., Gomes, L., Gavioli, E.C., Silva Junior, E.D.D., 2019. Fluoxetine and sertraline effects on rat distal cauda epididymis contraction, sperm count and sperm transit time through epididymis. *Eur J Pharmacol* 865, 172774.
- Bonney, R.C., Dixon, A.F., Fleming, D., 1979. Cyclic changes in the circulating and urinary levels of ovarian steroids in the adult female owl monkey (*Aotus trivirgatus*). *J Reprod Fertil* 56, 271-280.
- Brandt, N., Rune, GM, 2020. Sex-dependency of oestrogen-induced structural synaptic plasticity: Inhibition of aromatase versus application of estradiol in rodents. *Eur J Neurosci* 52, 2548-2559.
- Bromberger, J.T., Epperson, C.N., 2018. Depression During and After the Perimenopause: Impact of Hormones, Genetics, and Environmental Determinants of Disease. *Obstet Gynecol Clin North Am* 45, 663-678.
- Bryant, D.N., Sheldahl, L.C., Marriott, L.K., Shapiro, R.A., Dorsa, D.M., 2006. Multiple pathways transmit neuroprotective effects of gonadal steroids. *Endocrine* 29, 199-207.
- Channing, C.P., Fowler, S., Engel, B., Vitek, K., 1977. Failure of daily injections of ketamine HCL to adversely alter menstrual cycle length, blood estrogen, and progesterone levels in the rhesus monkey. *Proc Soc Exp Biol Med* 155, 615-619.
- Clayton, A.H., Warnock, J.K., Kornstein, S.G., Pinkerton, R., Sheldon-Keller, A., McGarvey, E.L., 2004. A placebo-controlled trial of bupropion SR as an antidote for selective serotonin reuptake inhibitor-induced sexual dysfunction. *J Clin Psychiatry* 65, 62-67.
- Dalla, C., Pitychoutis, P.M., Kokras, N., Papadopoulou-Daifoti, Z., 2010. Sex differences in animal models of depression and antidepressant response. *Basic Clin Pharmacol Toxicol* 106, 226-233.
- Damoiseaux, V.A., Proost, J.H., Jiawan, V.C., Melgert, B.N., 2014. Sex differences in the pharmacokinetics of antidepressants: influence of female sex hormones and oral contraceptives. *Clin Pharmacokinet* 53, 509-519.
- de Santi, F., Beltrame, F.L., Rodrigues, B.M., Junior, M., Scaramelle, N.F., Lopes, F.L., Cerri, P.S., Sasso-Cerri, E., 2020. Venlafaxine-induced damage to seminiferous epithelium, spermiogenesis, and sperm parameters in rats: A correlation with high estrogen levels. *Andrology*.
- Eid, A.H., Gad, A.M., Fikry, E.M., Arab, HH, 2019. Venlafaxine and carvedilol ameliorate testicular impairment and disrupted spermatogenesis in rheumatoid arthritis by targeting AMPK/ERK and PI3K/AKT/mTOR pathways. *Toxicol Appl Pharmacol* 364, 83-96.
- El Shehaby, D.M., El-Mahdy, R.I., Ahmed, A.M., Hosny, A., Abd El-Rady, N.M., 2020. Neurobehavioral, testicular and erectile impairments of chronic ketamine administration: Pathogenesis and ameliorating effect of N-acetyl cysteine. *Reprod Toxicol* 96, 57-66.
- ElMazoudy, R., El-Abd, K., Mekki, D., Kamel, K., 2020. Developmental effects on hypothalamic, hypophyseal, testicular and steroidogenic patterns of sertraline-exposed male rats by accumulated doses from juvenile to puberty. *Ecotoxicol Environ Saf* 188, 109840.

- Erdemir, F., Atilgan, D., Firat, F., Markoc, F., Parlaktas, B.S., Sogut, E., 2014. The effect of sertraline, paroxetine, fluoxetine and escitalopram on testicular tissue and oxidative stress parameters in rats. *Int Braz J Urol* 40, 100-108.
- Fooladi, E., Davis, S.R., 2012. An update on the pharmacological management of female sexual dysfunction. *Expert Opin Pharmacother* 13, 2131-2142.
- Frazer, A., 1997. Pharmacology of antidepressants. *J Clin Psychopharmacol* 17 Suppl 1, 2S-18S.
- Freeman, E.W., Sammel, M.D., Lin, H., Nelson, D.B., 2006. Associations of hormones and menopausal status with depressed mood in women with no history of depression. *Arch Gen Psychiatry* 63, 375-382.
- Garay, R.P., Zarate, C.A., Jr., Charpeaud, T., Citrome, L., Correll, C.U., Hameg, A., Llorca, P.M., 2017. Investigational drugs in recent clinical trials for treatment-resistant depression. *Expert Rev Neurother* 17, 593-609.
- Gould, E.M., 2008. The effect of ketamine/xylazine and carbon dioxide on plasma luteinizing hormone releasing hormone and testosterone concentrations in the male Norway rat. *Lab Anim* 42, 483-488.
- Gururajan, A., Reif, A., Cryan, J.F., Slattery, D.A., 2019. The future of rodent models in depression research. *Nat Rev Neurosci* 20, 686-701.
- Häggström, M., Richfield, D., 2014. Diagram of the pathways of human steroidogenesis. *WikiJournal of Medicine* 1.
- Hamdi, H., 2019. The preventive role of wheat germ oil against sertraline-induced testicular damage in male albino rats. *Andrologia* 51, e13369.
- Hermann, B., Vollmer, I., Holsboer, F., Rupprecht, R., 2001. Antidepressants do not modulate estrogen receptor alpha-mediated gene expression. *J Neural Transm (Vienna)* 108, 1197-1202.
- Hu, J., Zhang, Z., Shen, W.J., Azhar, S., 2010. Cellular cholesterol delivery, intracellular processing and utilization for biosynthesis of steroid hormones. *Nutr Metab (Lond)* 7, 47.
- Hutz, R.J., Dierschke, D.J., Wolf, R.C., 1988. Menstrual cycles in rhesus monkeys (*Macaca mulatta*) are unaffected by a single dose of the anesthetics ketamine and xylazine administered during the midfollicular phase at laparoscopy. *Am J Primatol* 15, 79-84.
- Ilgin, S., Aydogan-Kilic, G., Baysal, M., Kilic, V., Ardic, M., Ucarcan, S., Atli, O., 2018. Toxic Effects of Trazodone on Male Reproductive System via Disrupting Hypothalamic-Pituitary-Testicular Axis and Inducing Testicular Oxidative Stress. *Oxid Med Cell Longev* 2018, 7196142.
- Ilgin, S., Kilic, G., Baysal, M., Kilic, V., Korkut, B., Ucarcan, S., Atli, O., 2017. Citalopram Induces Reproductive Toxicity in Male Rats. *Birth Defects Res* 109, 475-485.
- Ishak, W.W., Bokarius, A., Jeffrey, J.K., Davis, M.C., Bakhta, Y., 2010. Disorders of orgasm in women: a literature review of etiology and current treatments. *J Sex Med* 7, 3254-3268.
- Islin, J., Munkboel, C.H., Styris, B., 2018. Steroidogenic disruptive effects of the serotonin-noradrenaline reuptake inhibitors duloxetine, venlafaxine and tramadol in the H295R cell assay and in a recombinant CYP17 assay. *Toxicol In Vitro* 47, 63-71.
- Kankash, A., Shariati, M., Khatamsaz, S., 2014. Effects of bupropion on physiology of reproduction in adult male rats. *J. Appl. Environ. Biol. Sci* 4, 106-110.
- Kilic, S., Ergin, H., Baydinc, Y.C., 2005. Venlafaxine extended release for the treatment of patients with premature ejaculation: a pilot, single-blind, placebo-controlled, fixed-dose crossover study on short-term administration of an antidepressant drug. *Int J Androl* 28, 47-52.
- Kingsberg, S.A., Woodard, T., 2015. Female sexual dysfunction: focus on low desire. *Obstet Gynecol* 125, 477-486.

- Kokras, N., Antoniou, K., Mikail, H.G., Kafetzopoulos, V., Papadopoulou-Daifoti, Z., Dalla, C., 2015. Forced swim test: What about females? *Neuropharmacology* 99, 408-421.
- Kokras, N., Dalla, C., 2014. Sex differences in animal models of psychiatric disorders. *Br J Pharmacol* 171, 4595-4619.
- Kokras, N., Dalla, C., 2017. Preclinical sex differences in depression and antidepressant response: Implications for clinical research. *J Neurosci Res* 95, 731-736.
- Kokras, N., Dalla, C., Papadopoulou-Daifoti, Z., 2011. Sex differences in pharmacokinetics of antidepressants. *Expert Opin Drug Metab Toxicol* 7, 213-226.
- Kokras, N., Dalla, C., Sideris, A.C., Dendi, A., Mikail, H.G., Antoniou, K., Papadopoulou-Daifoti, Z., 2012. Behavioral sexual dimorphism in models of anxiety and depression due to changes in HPA axis activity. *Neuropharmacology* 62, 436-445.
- Kokras, N., Hodes, G.E., Bangasser, D.A., Dalla, C., 2019. Sex differences in the hypothalamic-pituitary-adrenal axis: An obstacle to antidepressant drug development? *Br J Pharmacol* 176, 4090-4106.
- Kokras, N., Pastromas, N., Papasava, D., de Bournonville, C., Cornil, C.A., Dalla, C., 2018. Sex differences in behavioral and neurochemical effects of gonadectomy and aromatase inhibition in rats. *Psychoneuroendocrinology* 87, 93-107.
- Kokras, N., Polissidis, A., Antoniou, K., Dalla, C., 2017. Head shaking in the forced swim test: A robust but unexplored sex difference. *Pharmacol Biochem Behav* 152, 90-96.
- Kumar, A., Foster, T.C., 2020. G Protein-Coupled Estrogen Receptor: Rapid Effects on Hippocampal-Dependent Spatial Memory and Synaptic Plasticity. *Front Endocrinol (Lausanne)* 11, 385.
- Lazzara, R., Blazquez, M., Porte, C., Barata, C., 2012. Low environmental levels of fluoxetine induce spawning and changes in endogenous estradiol levels in the zebra mussel *Dreissena polymorpha*. *Aquat Toxicol* 106-107, 123-130.
- Lee, C.J., Do, B.R., Kim, J.K., Yoon, Y.D., 2000. Pentobarbital and ketamine suppress serum concentrations of sex hormones in the female rat. *J Anesth* 14, 187-190.
- Li, X., Wang, S., Zhang, L., Zhang, L., Liu, J., Luo, H., Gou, K., Cui, S., 2019. Amitriptyline plays important roles in modifying the ovarian morphology and improving its functions in rats with estradiol valerate-induced polycystic ovary. *Arch Pharm Res* 42, 344-358.
- Lister, A., Regan, C., Van Zwol, J., Van Der Kraak, G., 2009. Inhibition of egg production in zebrafish by fluoxetine and municipal effluents: a mechanistic evaluation. *Aquat Toxicol* 95, 320-329.
- Lupu, D., Pop, A., Cherfan, J., Kiss, B., Loghin, F., 2015. In vitro modulation of estrogen receptor activity by norfluoxetine. *Clujul Med* 88, 386-390.
- Lupu, D., Sjodin, M.O.D., Varshney, M., Lindberg, J., Loghin, F., Ruegg, J., 2017. Fluoxetine modulates sex steroid levels in vitro. *Clujul Med* 90, 420-424.
- Lupu, D., Varshney, M.K., Mucs, D., Inzunza, J., Norinder, U., Loghin, F., Nalvarte, I., Ruegg, J., 2018. Fluoxetine Affects Differentiation of Midbrain Dopaminergic Neurons In Vitro. *Mol Pharmacol* 94, 1220-1231.
- Malaivijitnond, S., Takenaka, O., Sankai, T., Yoshida, T., Cho, F., Yoshikawa, Y., 1998. Effects of single and multiple injections of ketamine hydrochloride on serum hormone concentrations in male cynomolgus monkeys. *Lab Anim Sci* 48, 270-274.
- McNeil, C.M., Murray, M., 1996. Inhibition of microsomal cytochromes P450 in rat liver by the tricyclic antidepressant drug desipramine and its primary oxidized metabolites. *Biochem Pharmacol* 51, 15-20.
- Meinhardt, W., Kropman, R.F., Vermeij, P., 1999. Comparative tolerability and efficacy of treatments for impotence. *Drug Saf* 20, 133-146.
- Mennigen, J.A., Lado, W.E., Zamora, J.M., Duarte-Guterman, P., Langlois, V.S., Metcalfe, C.D., Chang, J.P., Moon, T.W., Trudeau, V.L., 2010. Waterborne fluoxetine disrupts the

- reproductive axis in sexually mature male goldfish, *Carassius auratus*. *Aquat Toxicol* 100, 354-364.
- Mennigen, J.A., Martyniuk, C.J., Crump, K., Xiong, H., Zhao, E., Popescu, J., Anisman, H., Cossins, A.R., Xia, X., Trudeau, V.L., 2008. Effects of fluoxetine on the reproductive axis of female goldfish (*Carassius auratus*). *Physiol Genomics* 35, 273-282.
- Mennigen, J.A., Zamora, J.M., Chang, J.P., Trudeau, V.L., 2017. Endocrine disrupting effects of waterborne fluoxetine exposure on the reproductive axis of female goldfish, *Carassius auratus*. *Comp Biochem Physiol C Toxicol Pharmacol* 202, 70-78.
- Miller, M.K., Smith, J.R., Norman, J.J., Clayton, A.H., 2018. Expert opinion on existing and developing drugs to treat female sexual dysfunction. *Expert Opin Emerg Drugs* 23, 223-230.
- Miryala, C.S., Hiegel, C., Uphouse, L., 2013. Sprague-Dawley and Fischer female rats differ in acute effects of fluoxetine on sexual behavior. *J Sex Med* 10, 350-361.
- Monson, C., Young, G., Schultz, I., 2019. In vitro exposure of vitellogenic rainbow trout ovarian follicles to endocrine disrupting chemicals can alter basal estradiol-17 β production and responsiveness to a gonadotropin challenge. *Aquat Toxicol* 217, 105346.
- Montagnini, B.G., Bortolan, S., dos Santos, B.D., Moreno, A.P., Camin Nde, A., Gerardin, D.C., Moreira, E.G., 2013. Evaluation of escitalopram, sertraline, and methylphenidate in the immature rat uterotrophic assay. *Int J Toxicol* 32, 426-430.
- Munkboel, C.H., Larsen, L.W., Weisser, J.J., Mobjerg Kristensen, D., Styris have, B., 2018. Sertraline Suppresses Testis and Adrenal Steroid Production and Steroidogenic Gene Expression While Increasing LH in Plasma of Male Rats Resulting in Compensatory Hypogonadism. *Toxicol Sci* 163, 609-619.
- Nazian, S.J., 1988. Serum concentrations of reproductive hormones after administration of various anesthetics to immature and young adult male rats. *Proc Soc Exp Biol Med* 187, 482-487.
- Okano, M., Yazaki, T., Kawai, Y., Yamauchi, T., Nohara, Y., Furuya, I., Miwa, A., Kawara, M., Tateishi, T., 1988. [Application of the polyamide fiber to denture base. (Part II) Affection of surface treatment and incorporation of fibers to reinforce PMMA resin]. *Nichidai Koko Kagaku* 14, 46-53.
- Oyama, T., Toyota, M., Shinozaki, Y., Kudo, T., 1977. Effects of morphine and ketamine anaesthesia and surgery on plasma concentrations of luteinizing hormone, testosterone and cortisol in man. *Br J Anaesth* 49, 983-990.
- Pacher, P., Kohegyi, E., Kecskemeti, V., Furst, S., 2001. Current trends in the development of new antidepressants. *Curr Med Chem* 8, 89-100.
- Papakostas, G.I., Miller, K.K., Petersen, T., Sklar sky, K.G., Hilliker, S.E., Klibanski, A., Fava, M., 2006. Serum prolactin levels among outpatients with major depressive disorder during the acute phase of treatment with fluoxetine. *J Clin Psychiatry* 67, 952-957.
- Patel, K., Allen, S., Haque, M.N., Angelescu, I., Baumeister, D., Tracy, D.K., 2016. Bupropion: a systematic review and meta-analysis of effectiveness as an antidepressant. *Ther Adv Psychopharmacol* 6, 99-144.
- Pawluski, J.L., Kokras, N., Charlier, T.D., Dalla, C., 2020. Sex matters in neuroscience and neuropsychopharmacology. *Eur J Neurosci* 52, 2423-2428.
- Pirraglia, P.A., Stafford, R.S., Singer, D.E., 2003. Trends in prescribing of selective serotonin reuptake inhibitors and other newer antidepressant agents in adult primary care. *Primary care companion to the Journal of clinical psychiatry* 5, 153.
- Przegalinski, E., Warchol-Kania, A., Budziszewska, B., Jaworska, L., 1987. Effect of repeated administration of antidepressant drugs on the serum and brain concentration of testosterone and its metabolites. *Pol J Pharmacol Pharm* 39, 683-689.
- Puga-Olguin, A., Rodriguez-Landa, J.F., Rovirosa-Hernandez, M.J., German-Ponciano, L.J., Caba, M., Meza, E., Guillen-Ruiz, G., Olmos-Vazquez, O.J., 2019. Long-term ovariectomy

- increases anxiety- and despair-like behaviors associated with lower Fos immunoreactivity in the lateral septal nucleus in rats. *Behav Brain Res* 360, 185-195.
- Puri, C.P., Puri, V., Anand Kumar, T.C., 1981. Serum levels of testosterone, cortisol, prolactin and bioactive luteinizing hormone in adult male rhesus monkeys following cage-restraint or anaesthetizing with ketamine hydrochloride. *Acta Endocrinol (Copenh)* 97, 118-124.
- Pyke, R.E., 2020. Sexual Performance Anxiety. *Sex Med Rev* 8, 183-190.
- Qi, L., Liu, J.Y., Zhu, Y.L., Liu, W., Zhang, S.D., Liu, W.B., Jiang, J.J., 2017. Toxic effects of ketamine on reproductive system via disrupting hypothalamic-pituitary-testicular axis. *Eur Rev Med Pharmacol Sci* 21, 1967-1973.
- Rodas-Martinez, A.Z., Canales, D., Brousset, D.M., Swanson, W.F., Romano, MC, 2013. Assessment of adrenocortical and gonadal hormones in male spider monkeys (*Ateles geoffroyi*) following capture, restraint and anesthesia. *Zoo Biol* 32, 641-647.
- Ryan, J., Burger, H.G., Szoek, C., Lehert, P., Ancelin, M.L., Henderson, V.W., Dennerstein, L., 2009. A prospective study of the association between endogenous hormones and depressive symptoms in postmenopausal women. *Menopause* 16, 509-517.
- Saad, M.A., El-Sahar, A.E., Sayed, R.H., Elbaz, E.M., Helmy, H.S., Senousy, M.A., 2019. Venlafaxine Mitigates Depressive-Like Behavior in Ovariectomized Rats by Activating the EPO/EPOR/JAK2 Signaling Pathway and Increasing the Serum Estradiol Level. *Neurotherapeutics* 16, 404-415.
- Sakr, H.F., Abbas, A.M., Elsamanoudy, A.Z., Ghoneim, F.M., 2015. Effect of fluoxetine and resveratrol on testicular functions and oxidative stress in a rat model of chronic mild stress-induced depression. *J Physiol Pharmacol* 66, 515-527.
- Schlosser, R., Wetzel, H., Dorr, H., Rossbach, W., Hiemke, C., Benkert, O., 2000. Effects of subchronic paroxetine administration on night-time endocrinological profiles in healthy male volunteers. *Psychoneuroendocrinology* 25, 377-388.
- Sellers, K., Raval, P., Srivastava, D.P., 2015. Molecular signature of rapid estrogen regulation of synaptic connectivity and cognition. *Front Neuroendocrinol* 36, 72-89.
- Shansky, R.M., Woolley, C.S., 2016. Considering Sex as a Biological Variable Will Be Valuable for Neuroscience Research. *J Neurosci* 36, 11817-11822.
- Shebak, S.S., Varma, A., 2014. Low testosterone levels associated with venlafaxine use: a case report. *Prim Care Companion CNS Disord* 16.
- Silva de Assis, H.C., Simmons, D.B., Zamora, J.M., Lado, W.E., Al-Ansari, A.M., Sherry, J.P., Blais, J.M., Metcalfe, C.D., Trudeau, V.L., 2013. Estrogen-like effects in male goldfish co-exposed to fluoxetine and 17 alpha-ethinylestradiol. *Environ Sci Technol* 47, 5372-5382.
- Soares, C.N., Almeida, O.P., Joffe, H., Cohen, L.S., 2001. Efficacy of estradiol for the treatment of depressive disorders in perimenopausal women: a double-blind, randomized, placebo-controlled trial. *Arch Gen Psychiatry* 58, 529-534.
- Sonntag, A., Rothe, B., Guldner, J., Yassouridis, A., Holsboer, F., Steiger, A., 1996. Trimipramine and imipramine exert different effects on the sleep EEG and on nocturnal hormone secretion during treatment of major depression. *Depression* 4, 1-13.
- Sramek, J.J., Cutler, N.R., 2011. The impact of gender on antidepressants. *Curr Top Behav Neurosci* 8, 231-249.
- Sramek, J.J., Murphy, M.F., Cutler, N.R., 2016. Sex differences in the psychopharmacological treatment of depression. *Dialogues Clin Neurosci* 18, 447-457.
- Steiger, A., von Bardeleben, U., Wiedemann, K., Holsboer, F., 1991. Sleep EEG and nocturnal secretion of testosterone and cortisol in patients with major endogenous depression during acute phase and after remission. *J Psychiatr Res* 25, 169-177.

- Sterzik, K., Nitsch, C.D., Korda, P., Sasse, V., Rosenbusch, B., Marx, T., Traub, E., 1994. [The effect of different anesthetic procedures on hormone levels in women. Studies during an in vitro fertilization-embryo transfer (IVF-ET) program]. *Anaesthesist* 43, 738-742.
- Taxier, L.R., Gross, K.S., Frick, K.M., 2020. Oestradiol as a neuromodulator of learning and memory. *Nat Rev Neurosci* 21, 535-550.
- Taylor, G.T., Farr, S., Klinga, K., Weiss, J., 2004. Chronic fluoxetine suppresses circulating estrogen and the enhanced spatial learning of estrogen-treated ovariectomized rats. *Psychoneuroendocrinology* 29, 1241-1249.
- Trickler, W.J., Guo, X., Cuevas, E., Ali, S.F., Paule, M.G., Kanungo, J., 2014. Ketamine attenuates cytochrome p450 aromatase gene expression and estradiol-17beta levels in zebrafish early life stages. *J Appl Toxicol* 34, 480-488.
- Uphouse, L., Hensler, J.G., Sarkar, J., Grossie, B., 2006. Fluoxetine disrupts food intake and estrous cyclicity in Fischer female rats. *Brain Res* 1072, 79-90.
- Vitezic, D., Pelcic, J.M., 2002. Erectile dysfunction: oral pharmacotherapy options. *Int J Clin Pharmacol Ther* 40, 393-403.
- Weissheimer, K.V., Herod, S.M., Cameron, J.L., Bethea, C.L., 2010. Interactions of corticotropin-releasing factor, urocortin and citalopram in a primate model of stress-induced amenorrhea. *Neuroendocrinology* 92, 224-234.
- Westlund Tam, L., Parry, B.L., 2003. Does estrogen enhance the antidepressant effects of fluoxetine? *J Affect Disord* 77, 87-92.
- Williams, A.V., Trainor, B.C., 2018. The impact of sex as a biological variable in the search for novel antidepressants. *Front Neuroendocrinol* 50, 107-117.
- Yamazaki, H., Shaw, P.M., Guengerich, F.P., Shimada, T., 1998. Roles of cytochromes P450 1A2 and 3A4 in the oxidation of estradiol and estrone in human liver microsomes. *Chem Res Toxicol* 11, 659-665.
- Yardimci, A., Ulker, N., Bulmus, O., Kaya, N., Colakoglu, N., Ozcan, M., Canpolat, S., Kelestimur, H., 2019. Effects of long-term paroxetine or bupropion treatment on puberty onset, reproductive and feeding parameters in adolescent male rats. *Andrologia* 51, e13268.
- Yohn, C.N., Gergues, M.M., Samuels, B.A., 2017. The role of 5-HT receptors in depression. *Mol Brain* 10, 28.
- Zaidi, P., Wickings, E.J., Nieschlag, E., 1982. The effects of ketamine HCl and barbiturate anaesthesia on the metabolic clearance and production rates of testosterone in the male rhesus monkey, *Macaca mulatta*. *J Steroid Biochem* 16, 463-466.
- Zarembka, F.R., Koller, D.E., Plotka, E.D., 1989. Effect of ether or ketamine anesthesia on rat uterine estrogen and progesterone receptors. *Clin Chem* 35, 143-145.
- Zheng, Y., Wu, M., Gao, T., Meng, L., Ding, X., Meng, Y., Jiao, Y., Luo, P., He, Z., Sun, T., Zhang, G., Shi, X., Rong, W., 2020. GPER-Deficient Rats Exhibit Lower Serum Corticosterone Level and Increased Anxiety-Like Behavior. *Neural Plast* 2020, 8866187.

Figure 1: Biosynthesis of steroid hormones. Image created with BioRender.com based on an image from ([Häggström and Richfield, 2014](#))

Journal Pre-proof

Figure 2: Testosterone levels in adult male and female Wistar rats. An asterisk (*) denotes a significant difference, as evidenced by the ANOVA analysis, between sertraline-treated rats and the corresponding vehicle-treated group of the same sex. A cross sign (+) indicates a significant sex difference between females and the corresponding male group. N = 7–9 per group, Means \pm Standard Error of Mean. Adapted under licence from ([Kokras et al., 2015](#)).

Authors	Drug	Sex	Strain	Age (weeks)	Dose (mg/kg)	Duration (days)	Serum	Brain
Ilgin <i>et al.</i> 2017	Citalopram	Male	W	12	5, 10, 20	28	↑	
Erdemir <i>et al.</i> 2014	Citalopram	Male	W, AI	22-24	10	60	↓	
Przegalinski <i>et al.</i> 1987	Citalopram	Male	W	Adult	10	21	↑	↑
Munkboel <i>et al.</i> 2018	Sertraline	Male	SD	9	1.25, 5, 20	15	↔	↓
Erdemir <i>et al.</i> 2014	Sertraline	Male	W, AI	22-24	10	60	↓	
Atli <i>et al.</i> 2017	Sertraline	Male	W	8-10	20	30	↑	
Hamdi 2019	Sertraline	Male	W	7-9	15.63	28	↑	
ElMazoudy <i>et al.</i> 2020	Sertraline	Male	W	3	12	28	↓	
Bezerra <i>et al.</i> 2019	Sertraline	Male	W	8-12	20	21	↓	
Kokras <i>et al.</i> 2015	Sertraline	Male	W	Adult	10, 40	3	↓	
Kokras <i>et al.</i> 2015	Sertraline	Female	W	Adult	10, 40	3	↔	
Allouh <i>et al.</i> 2014	Paroxetine	Male	SD, AI	Adult	10	20	↑	
Erdemir <i>et al.</i> 2014	Paroxetine	Male	W, AI	22-24	10	60	↓	
Mennigen <i>et al.</i> 2010	Fluoxetine	Male	G		54 (µg/L)		↓	
Ayala <i>et al.</i> 2018	Fluoxetine	Male	CII	4-8	5	23	↓	
Erdemir <i>et al.</i> 2014	Fluoxetine	Male	W, AI	22-24	10	60	↔	
Bataineh and Daradka 2007	Fluoxetine	Male	SD	Adult	200	60	↓	
Sakr <i>et al.</i> 2015	Fluoxetine	Male	SD	young	10	28	↓	
Bezerra <i>et al.</i> 2019	Fluoxetine	Male	W	8-12	20	21	↓	
de Santi <i>et al.</i> 2020	Venlafaxine	Male	H	12	30	35, 65	↑	
Eid <i>et al.</i> 2019	Venlafaxine	Male	SD	Adult	50	20	↑	

Journal Pre-proof								
Przegalinski <i>et al.</i> 1987	Amitriptyline	Male	W	Adult	10	21	↓	↓
Li <i>et al.</i> 2019	Amitriptyline	Female	W	Adult	10	30	↔	
Przegalinski <i>et al.</i> 1987	Imipramine	Male	W	Adult	10	21	↓	↓
Yardimci <i>et al.</i> 2019	Bupropion	Male	SD	3	17	69	↑	
Kankash <i>et al.</i> 2014	Bupropion	Male	W				↑	
Hrushesky <i>et al.</i> 1988	Trazodone	Male	SD		5, 10, 20	28	↑	
Ashraf <i>et al.</i> 2014	Trazodone	Male	W	Adult	90	28	↓	
Trickler <i>et al.</i> 2014	Esketamine		Z	Embryos			↑	
Qi <i>et al.</i> 2017	Esketamine	Male	SD	5-6	20, 40, 60	21	↓	
Nazian 1988	Esketamine	Male	SD	3	200	22	↓	
Gould 2008	Esketamine	Male	SD	12	37.5	Single	↔	
El Shehaby <i>et al.</i> 2020	Esketamine	Male	AI	5-6	10	84	↓	
Lee <i>et al.</i> 2000	Esketamine	Female	SD	12	229	Acute	↓	
Puri <i>et al.</i> 1981	Esketamine	Male	RM	Adult	10	Single	↓	
Zaidi <i>et al.</i> 1982	Esketamine	Male	RM	Adult			↔	
Malaivijitnond <i>et al.</i> 1998	Esketamine	Male	CM	Adult	10	28	↔	
Rodas-Martinez <i>et al.</i> 2013	Esketamine	Male	SM	8-21 (yo)	10	Acute	↓	

Table 1: Serum or brain testosterone levels after the administration of antidepressants in published preclinical studies. The sex, strain, and the age of the animals (in weeks, or months) are indicated, when available. Variations in treatment doses, route of administration, and duration are indicated, when available. \uparrow : indicates increased testosterone levels after the treatment, \downarrow : indicates decreases testosterone levels after the treatment, and \Leftrightarrow : no changes in testosterone levels after the treatment. Strains: W = Wistar, Al = Albino, SD = Sprague-Dawley, G = Goldfish, CII = CII-Z, LE = Long-Evans, H = Holtzman, RM = Rhesus monkeys, CM = Cynomolgus monkeys, SM = Spider monkeys, Z = Zebrafish (Danio rerio). Age: yo = years old.

Authors	Drug	Gender	Status	Mean Age (years)	Dose (mg/day)	Duration (weeks)	Testosterone
Schlosser <i>et al.</i> 2000	Paroxetine	Men	Healthy	27	30	4	↔
Papakostas <i>et al.</i> 2006	Fluoxetine	Men	MDD			12	↔
Bell <i>et al.</i> 2006	Fluoxetine	Men	Depression	22	10 to 20	4	↔
Bell and Shipman 2000	Venlafaxine	Men	Depression	26	37.5-150	52	↓
Kilic <i>et al.</i> 2005	Venlafaxine	Men	Healthy	34	75	2	↔
Steiger <i>et al.</i> 1991	Amitriptyline	Men	MDD	46	150		↑
Sonntag <i>et al.</i> 1996	Imipramine	Men	MDD	43	5 to 150	28	↑
Oyama <i>et al.</i> 1977	Ketamine	Men	Healthy	32 and 36	2	Single	↓

Table 2: Testosterone levels after the administration of antidepressants in published clinical studies. The gender, health status, and the age of the participants (mean age) are indicated, when available. Variations in treatment doses and duration are indicated, when available. ↑: indicates increased testosterone levels after the treatment, ↓: indicates decreases testosterone levels after the treatment, and ↔: no changes in testosterone levels after the treatment. Status: MDD = Major Depressive Disorder.

Authors	Drug	Sex	Strain	Age (weeks)	Dose (mg/kg)	Duration (days)	Serum	Brain
Przegalinski <i>et al.</i> 1987	Citalopram	Male	W	Adult	10	21	↔	↔
Weissheimer <i>et al.</i> 2010	Citalopram	Female	CM		1.2-4.8	105	↓	
Kokras <i>et al.</i> 2015	Sertraline	Male	W	Adult	10, 40	3	↔	
Kokras <i>et al.</i> 2015	Sertraline	Female	W	Adult	10, 40	3	↔	
Pressley and Branas 2018	Fluoxetine	Male	G				↑	
Silva de Assis <i>et al.</i> 2013	Fluoxetine	Male	G				↑	
Mennigen <i>et al.</i> 2008	Fluoxetine	Female					↓	
Lister <i>et al.</i> 2009	Fluoxetine		Z				↓	
Kim and Hancock 1989	Fluoxetine	Female	MD				↓	
Monson <i>et al.</i> 2019	Fluoxetine	Female	RT				↔	
Lazzara <i>et al.</i> 2012	Fluoxetine	Male	ZM				↑	
Lazzara <i>et al.</i> 2012	Fluoxetine	Female	ZM				↑	
Uphouse <i>et al.</i> 2006	Fluoxetine	Female	F	Adult	10	12-23	↔	
Taylor <i>et al.</i> 2004	Fluoxetine	Female	LE		0.5-5		↓	
de Santi <i>et al.</i> 2020	Venlafaxine	Male	H	12	30	35, 65	↓	
Saad <i>et al.</i> 2019	Venlafaxine	Female	W	50	10		↑	
Przegalinski <i>et al.</i> 1987	Amitriptyline	Male	W	Adult	10	21	↔	↔
Li <i>et al.</i> 2019	Amitriptyline	Female	W	Adult	10	30	↑	
Przegalinski <i>et al.</i> 1987	Imipramine	Male	W	Adult	10	21	↔	↔
El-Khatib <i>et al.</i> 2020	Agomelatine	Female	W	12	40	10	↑	
Trickler <i>et al.</i> 2014	Esketamine		Z	Embryos			↓	

Journal Pre-proof							
Lee <i>et al.</i> 2000	Esketamine	Female	SD	12	229	Acute	↓
Bonney <i>et al.</i> 1979	Esketamine	Female	OM	Adult	10	Acute	↔
Channing <i>et al.</i> 1977	Esketamine	Female	RM	Adult	8-10	365	↔
Hutz <i>et al.</i> 1988	Esketamine	Female	RM	56-18 (yo)	15	Single	↔

Table 3: Serum or brain estrogen levels after the administration of antidepressants in published preclinical studies. The sex, strain, and the age of the animals (in weeks, or months) are indicated, when available. Variations in treatment doses, route of administration, and duration are indicated, when available. ↑: indicates increased estrogen levels after the treatment, ↓: indicates decreases estrogen levels after the treatment, and ↔: no changes in estrogen levels after the treatment. Strains: W = Wistar, Al = Albino, SD = Sprague-Dawley, G = Goldfish, CII = CII-Z, LE = Long-Evans, H = Holtzman, RM = Rhesus monkeys, CM= Cynomolgus monkeys, SM = Spider monkeys, Z = Zebrafish (*Danio rerio*). Age: yo = years old.

Authors	Drug	Gender	Status	Mean Age (years)	Dose (mg/day)	Duration (weeks)	Estrogen
Bao <i>et al.</i> 2004	Fluoxetine	Women	MDD	33	20	12	↑
Sterzik <i>et al.</i> 1994	Ketamine	Women	Healthy				↔

Table 4: Estrogen levels after the administration of antidepressants in published clinical studies. The gender, health status, and the age of the participants (mean age) are indicated, when available. Variations in treatment doses and duration are indicated, when available. ↑: indicates increased estrogen levels after the treatment, ↔: no changes in estrogen levels after the treatment. Status: MDD = Major Depressive Disorder.



